

ISSN 2311-4673
Journal of Pharmacy and Pharmaceutical Sciences
(Volume 2, Issue 2, 2014)

What is Cancer? What Causes Cancer?

Muhammad Shakil Suakeh Siddiqui^{1}, Ghulam Sarwar¹*

¹Faculty of Pharmacy, Jinnah University for Women, Karachi

INTRODUCTION

Cancer is a class of diseases characterized by out-of-control cell growth. There are over 100 different types of cancer, and each is classified by the type of cell that is initially affected.

Cancer harms the body when damaged cells divide uncontrollably to form lumps or masses of tissue called tumors (except in the case of leukemia where cancer prohibits normal blood function by abnormal cell division in the blood stream). Tumors can grow and interfere with the digestive, nervous, and circulatory systems, and they can release hormones that alter body function. Tumors that stay in one spot and demonstrate limited growth are generally considered to be benign.

More dangerous, or malignant, tumors form when two things occur:

1. a cancerous cell manages to move throughout the body using the blood or lymph systems, destroying healthy tissue in a process called invasion
2. that cell manages to divide and grow, making new blood vessels to feed itself in a process called angiogenesis.

When a tumor successfully spreads to other parts of the body and grows, invading and destroying other healthy tissues, it is said to have metastasized. This

*Corresponding author: mshakil@hotmail.com

process itself is called metastasis, and the result is a serious condition that is very difficult to treat.

How cancer spreads?

Scientists reported in Nature Communications (October 2012 issue)² that they have discovered an important clue as to why cancer cells spread. It has something to do with their adhesion (stickiness) properties. Certain molecular interactions between cells and the scaffolding that holds them in place (extracellular matrix) cause them to become unstuck at the original tumor site, they become dislodged, move on and then reattach themselves at a new site. The researchers say this discovery is important because cancer mortality is mainly due to metastatic tumors, those that grow from cells that have traveled from their original site to another part of the body. Only 10% of cancer deaths are caused by the primary tumors.

The scientists, from the Massachusetts Institute of Technology, say that finding a way to stop cancer cells from sticking to new sites could interfere with metastatic disease, and halt the growth of secondary tumors.

In 2007, cancer claimed the lives of about 7.6 million people in the world. Physicians and researchers who specialize in the study, diagnosis, treatment, and prevention of cancer are called oncologists.

What causes cancer?

Cancer is ultimately the result of cells that uncontrollably grow and do not die. Normal cells in the body follow an orderly path of growth, division, and death. Programmed cell death is called apoptosis, and when this process breaks down, cancer begins to form. Unlike regular cells, cancer cells do not experience programmed death and instead continue to grow and divide. This leads to a mass of abnormal cells that grows out of control.

Genes - the DNA type

Cells can experience uncontrolled growth if there are damages or mutations to DNA, and therefore, damage to the genes involved in cell division. Four key types of gene are responsible for the cell division process: oncogenes tell cells when to divide, tumor suppressor genes tell cells when not to divide, suicide genes control apoptosis and tell the cell to kill itself if something goes wrong, and DNA-repair genes instruct a cell to repair damaged DNA. Cancer occurs when a cell's gene mutations make the cell unable to correct DNA damage and unable to commit suicide. Similarly, cancer is a result of mutations that inhibit oncogene and tumor suppressor gene function, leading to uncontrollable cell growth.

Carcinogens

Carcinogens are a class of substances that are directly responsible for damaging DNA, promoting or aiding cancer. Tobacco, asbestos, arsenic, radiation such as gamma and x-rays, the sun, and compounds in car exhaust fumes are all examples of carcinogens. When our bodies are exposed to carcinogens, free radicals are formed that try to steal electrons from other molecules in the body. These free radicals damage cells and affect their ability to function normally.

Genes - the family type

Cancer can be the result of a genetic predisposition that is inherited from family members. It is possible

to be born with certain genetic mutations or a fault in a gene that makes one statistically more likely to develop cancer later in life.

What are the symptoms of cancer?

Cancer symptoms are quite varied and depend on where the cancer is located, where it has spread, and how big the tumor is. Some cancers can be felt or seen through the skin - a lump on the breast or testicle can be an indicator of cancer in those locations. Skin cancer (melanoma) is often noted by a change in a wart or mole on the skin. Some oral cancers present white patches inside the mouth or white spots on the tongue.

Other cancers have symptoms that are less physically apparent. Some brain tumors tend to present symptoms early in the disease as they affect important cognitive functions. Pancreas cancers are usually too small to cause symptoms until they cause pain by pushing against nearby nerves or interfere with liver function to cause a yellowing of the skin and eyes called jaundice. Symptoms also can be created as a tumor grows and pushes against organs and blood vessels. For example, colon cancers lead to symptoms such as constipation, diarrhea, and changes in stool size. Bladder or prostate cancers cause changes in bladder function such as more frequent or infrequent urination.

As cancer cells use the body's energy and interfere with normal hormone function, it is possible to present symptoms such as fever, fatigue, excessive sweating, anemia, and unexplained weight loss. However, these symptoms are common in several other maladies as well. For example, coughing and hoarseness can point to lung or throat cancer as well as several other conditions.

When cancer spreads, or metastasizes, additional symptoms can present themselves in the newly affected area. Swollen or enlarged lymph nodes are common and likely to be present early. If cancer spreads to the brain, patients may experience vertigo, headaches, or seizures. Spreading to the lungs may

cause coughing and shortness of breath. In addition, the liver may become enlarged and cause jaundice and bones can become painful, brittle, and break easily. Symptoms of metastasis ultimately depend on the location to which the cancer has spread.

How is cancer classified?

There are five broad groups that are used to classify cancer.

1. Carcinomas are characterized by cells that cover internal and external parts of the body such as lung, breast, and colon cancer.
2. Sarcomas are characterized by cells that are located in bone, cartilage, fat, connective tissue, muscle, and other supportive tissues.
3. Lymphomas are cancers that begin in the lymph nodes and immune system tissues.
4. Leukemias are cancers that begin in the bone marrow and often accumulate in the bloodstream.
5. Adenomas are cancers that arise in the thyroid, the pituitary gland, the adrenal gland, and other glandular tissues.

Cancers are often referred to by terms that contain a prefix related to the cell type in which the cancer originated and a suffix such as -sarcoma, -carcinoma, or just -oma. Common prefixes include:

- Adeno- = gland
- Chondro- = cartilage
- Erythro- = red blood cell
- Hemangio- = blood vessels
- Hepato- = liver
- Lipo- = fat
- Lympho- = white blood cell
- Melano- = pigment cell
- Myelo- = bone marrow
- Myo- = muscle
- Osteo- = bone
- Uro- = bladder

- Retino- = eye
- Neuro- = brain

How is cancer diagnosed and staged?

Early detection of cancer can greatly improve the odds of successful treatment and survival. Physicians use information from symptoms and several other procedures to diagnose cancer. Imaging techniques such as X-rays, CT scans, MRI scans, PET scans, and ultrasound scans are used regularly in order to detect where a tumor is located and what organs may be affected by it. Doctors may also conduct an endoscopy, which is a procedure that uses a thin tube with a camera and light at one end, to look for abnormalities inside the body.



Extracting cancer cells and looking at them under a microscope is the only absolute way to diagnose cancer. This procedure is called a biopsy. Other types of molecular diagnostic tests are frequently employed as well. Physicians will analyze your body's sugars, fats, proteins, and DNA at the molecular level. For example, cancerous prostate cells release a higher level of a chemical called PSA (prostate-specific antigen) into the bloodstream that can be detected by a blood test. Molecular diagnostics, biopsies, and imaging techniques are all used together to diagnose cancer.

How is cancer treated?

Cancer treatment depends on the type of cancer, the stage of the cancer (how much it has spread),

age, health status, and additional personal characteristics. There is no single treatment for cancer, and patients often receive a combination of therapies and palliative care. Treatments usually fall into one of the following categories: surgery, radiation, chemotherapy, immunotherapy, hormone therapy, or gene therapy.

Surgery

Surgery is the oldest known treatment for cancer. If a cancer has not metastasized, it is possible to completely cure a patient by surgically removing the cancer from the body. This is often seen in the removal of the prostate or a breast or testicle. After the disease has spread, however, it is nearly impossible to remove all of the cancer cells. Surgery may also be instrumental in helping to control symptoms such as bowel obstruction or spinal cord compression.

Radiation



Radiation treatment, also known as radiotherapy, destroys cancer by focusing high-energy rays on the cancer cells. This causes damage to the molecules that make up the cancer cells and leads them to commit suicide. Radiotherapy utilizes high-energy gamma-rays that are emitted from metals such as radium or high-energy x-rays that are created in a special machine. Early radiation treatments caused severe side-effects because the energy beams would damage normal, healthy tissue, but technologies have improved so that beams can be more accurately

targeted. Radiotherapy is used as a standalone treatment to shrink a tumor or destroy cancer cells (including those associated with leukemia and lymphoma), and it is also used in combination with other cancer treatments.

Chemotherapy

Chemotherapy⁴ utilizes chemicals that interfere with the cell division process - damaging proteins or DNA - so that cancer cells will commit suicide. These treatments target any rapidly dividing cells (not necessarily just cancer cells), but normal cells usually can recover from any chemical-induced damage while cancer cells cannot. Chemotherapy is generally used to treat cancer that has spread or metastasized because the medicines travel throughout the entire body. It is a necessary treatment for some forms of leukemia and lymphoma. Chemotherapy treatment occurs in cycles so the body has time to heal between doses. However, there are still common side effects such as hair loss, nausea, fatigue, and vomiting. Combination therapies often include multiple types of chemotherapy or chemotherapy combined with other treatment options.

Immunotherapy

Immunotherapy aims to get the body's immune system to fight the tumor. Local immunotherapy injects a treatment into an affected area, for example, to cause inflammation that causes a tumor to shrink. Systemic immunotherapy treats the whole body by administering an agent such as the protein interferon alpha that can shrink tumors. Immunotherapy can also be considered non-specific if it improves cancer-fighting abilities by stimulating the entire immune system, and it can be considered targeted if the treatment specifically tells the immune system to destroy cancer cells. These therapies are relatively young, but researchers have had success with treatments that introduce antibodies to the body that inhibit the growth of breast cancer cells. Bone marrow transplantation (hematopoietic stem cell transplantation) can also be considered

immunotherapy because the donor's immune cells will often attack the tumor or cancer cells that are present in the host.

Hormone therapy

Several cancers have been linked to some types of hormones, most notably breast and prostate cancer. Hormone therapy⁵ is designed to alter hormone production in the body so that cancer cells stop growing or are killed completely. Breast cancer hormone therapies often focus on reducing estrogen levels (a common drug for this is tamoxifen) and prostate cancer hormone therapies often focus on reducing testosterone levels. In addition, some leukemia and lymphoma cases can be treated with the hormone cortisone.

Gene therapy

The goal of gene therapy is to replace damaged genes with ones that work to address a root cause of cancer: damage to DNA. For example, researchers are trying to replace the damaged gene that signals cells to stop dividing (the p53 gene) with a copy of a working gene. Other gene-based therapies focus on further damaging cancer cell DNA to the point where the cell commits suicide. Gene therapy is a very young field and has not yet resulted in any successful treatments.

REFERENCES

1. Nature 2012. "How many different types of cancer are there? Cancer Research UK : Cancer Help UK". Retrieved 11 May (2012).
3. Anand P, Kunnumakkara AB, Kunnumakara AB, Sundaram C, Harikumar KB, Tharakan ST, Lai OS, Sung B, Aggarwal BB. "Cancer is a preventable disease that requires major lifestyle changes". *Pharm. Res.*25 (9): 2097–116 (2008). doi:10.1007/ s11095-008-9661-9.PMC 2515569.PMID 18626751.

4. Kinzler, Kenneth W.; Vogelstein, Bert. "Introduction".*The genetic basis of human cancer* (2nd, illustrated, revised ed.). New York: McGraw-Hill, Medical Pub. Division. p. 5.ISBN 978-0-07-137050-9 (2002).
5. Jemal A, Bray, F, Center, MM, Ferlay, J, Ward, E, Forman, D. "Global cancer statistics". *CA: a cancer journal for clinicians* 61 (2): 69–90. doi:10.3322/caac. 20107. PMID 21296855 (2011).